FOUNDATION COURSE EXAMINATION

June 2013

PAPER 4: Business Mathematics and Statistics Fundamentals

Full Marks: 50

Answer all the questions. Each question carries 1 mark. Each Question has four alternatives. Indicate the correct alternative by darkening the circle on the OMR Sheet.

- 1. If x : y = 3 : 4, y : z = 5 : 6 and z : w = 7 : 8, then x : y : z : w is
 - (a) 3:5:7:8
 - (b) 3:4:6:8
 - (c) 105: 140: 168: 192
 - (d) 3:4:24:192
- 2. The average monthly consumption of petrol for a car for 12 months is 160 litre, If the average monthly consumption for first 8 months is 145 litre, then the average monthly consumption of petrol for the last 4 months is
 - (a) 190 litre
 - (b) 165 litre
 - (c) 180 litre
 - (d) 175 litre
- If 9 men working 10 hours daily can complete a job in 10 days, then 15 men working 6 hours daily shall complete same job in
 - (a) 6 days
 - (b) 8 days
 - (c) 10 days
 - (d) none of the above
- 4. In 25 years at 8% p.a. simple interest, a sum becomes ₹4629. The amount of sum is
 - (a) ₹1534
 - (b) ₹1453
 - (c) ₹1435
 - (d) ₹1543
- 5. True discount at interest rate 5% p.a. is ₹40. A bill of ₹1240 is due in
 - (a) 6 months
 - (b) 8 months
 - (c) 1 year
 - (d) $1\frac{1}{2}$ year

- 6. If $3^{x+1} + 3^{x-1} = 270$, then the value of x is
 - (a) 5
 - (b) 3
 - (c) 4
 - (d) 8
- 7. If $x = 7 4\sqrt{3}$, then the value of $\sqrt{\chi} + \frac{1}{\sqrt{\chi}}$ is
 - (a) $2\sqrt{3}$
 - (b) $4\sqrt{3}$
 - (c) 4
 - (d) 6
- 8. If x = 2 + 3i then $x^2 4x + 13is$
 - (a) 0
 - (b) 1
 - (c) 2
 - (d) none of the above
- 9. Value of log 144 with base $2\sqrt{3}$ is
 - (a) 2
 - (b) 3
 - (c) 4
 - (d) none of the above
- 10. If P and Q be two non-empty sets, then $(P-Q)\cap(Q-P)\cap(P\cap Q)$
 - (a) Φ
 - (b) P n Q
 - (c) PuQ
 - (d) none of the above
- 11. The ratio of $({}^{2n}P_n + {}^{2n}C_n)$: $({}^{2n}P_n {}^{2n}C_n)$ is
 - (a) (n+1): (n-1)
 - (b) (n-1): (n+1)
 - (c) (n! + 1):(n!-1)
 - (d) (n+1)!:(n-1)!

- 12. A quantity p varies directly as t and another quantity q varies inversely as t. When t=2, p+q=1 and when t=3, p+q=8. When t=4, p+q=1
 - (a) 8.5
 - (b) 9
 - (c) 9.5
 - (d) 8
- 13. If p and q be respectively two statements "he is tall" and "he is intelligent" then the symbolic form of the statement "it is not true that he is short or dull" is
 - (a) ~ p ^ ~q
 - (b) $\sim (\sim p \vee q)$
 - (c) ~(~p,~q)
 - (d) \sim (p $_{\vee}\sim$ q)
- 14. The area of the triangle with sides of 9 cm, 12 cm and 15 cm is
 - (a) 50 sq cm
 - (b) 54 sq cm
 - (c) 56 sq cm
 - (d) none of the above
- 15. Sum of all interior angles of a decagon is
 - (a) 1200°
 - (b) 1300°
 - (c) 1400°
 - (d) none of the above
- 16. The circumference of the base of a cylinder is 66 cm. If the volume of the cylinder be 3465 cu cm, then the height of the cylinder is
 - (a) 10 cm
 - (b) 10.5 cm
 - (c) 11 cm
 - (d) 12.5 cm
- 17. Base of a right prism is an equilateral triangle of side 4 cm. If the volume of the prism is $60\sqrt{3}$ cu cm, then its height is
 - (a) 20 cm
 - (b) 12 cm
 - (c) 18 cm
 - (d) 15 cm
- 18. Three solid spherical balls of diameters 10, 8 and 6 cm respectively are melted into one solid ball. Its diameter is
 - (a) 11 cm
 - (b) 12 cm
 - (c) 10 cm
 - (d) 13 cm

- 19. Diameter of the base of a conical tent is 14 metre and its height is 9 metre. Its volume is
 - (a) 450 cu metre
 - (b) 460 cu metre
 - (c) 470 cu metre
 - (d) none of the above
- 20. A right pyramid of height 4 cm stands on a square base of side 3 cm. Its volume is
 - (a) 10 cu cm
 - (b) 12 cu cm
 - (c) 14 cu cm
 - (d) none of the above
- 21. The perimeter of a rhombus is 64 cm and one of the diagonals is 20 cm. The area of the rhombus is
 - (a) 80√39sq cm
 - (b) 20√39sqcm
 - (c) 40√39sqcm
 - (d) 39√40 sq cm
- 22. If the point R divides the line segment joining (6, 3) and (9, 12) internally in the ratio 2:1, then R is
 - (a) (5,6)
 - (b) (6,7)
 - (c) (7,8)
 - (d) none of the above
- 23. The perpendicular distance of the point $\left(\frac{3}{2}, -\frac{5}{2}\right)$ from

the line 2x - y = 3 is

- (a) $\frac{2}{\sqrt{5}}$ units
- (b) √5 units
- (c) $\frac{1}{2}$ units
- (d) $\frac{\sqrt{5}}{2}$ units
- 24. If the centre of the cricle $x^2 + y^2 + 6x 3by + 2 = 0$ be (-3, 6), then the value of b is
 - (a) 3
 - (b) 4
 - (c) -3
 - (d) -4

- 25. The eccentricity of an ellipse is $\frac{\sqrt{3}}{2}$ units and major axis is
 - 12 units. The minor axis is
 - (a) 4 units
 - (b) 2 units
 - (c) 6 units
 - (d) $2\sqrt{3}$ units
- 26. The vertex of the parabola $x^2 + 8x + 12y + 4 = 0$ is
 - (a) (-4,1)
 - (b) (4, -1)
 - (c) (-4,-1)
 - (d) none of the above
- 27. $\lim_{X \to 0} \frac{x + \log(1 + x)}{e^{X} 1} =$
 - (a) 1
 - (b) 4
 - (c) 1.5
 - (d) none of the above
- 28. If $f(x) = \frac{3x+2}{2x+3}$, then $f(-x)f(-\frac{1}{x})$ is
 - (a) -1
 - (b) 1
 - (c) $\frac{2}{3}$
 - (d) $\frac{3}{2}$
- 29. The derivative of \sqrt{x} with respect to log x is
 - (a) √x
 - (b) 2√x
 - (c) X
 - (d) $\frac{\sqrt{x}}{2}$
- 30. A firm produces x units of output per week at a total cost of $\frac{1}{3}$ x³ 7x² + 45x. The level of output at which the
 - marginal cost attains the minimum is
 - (a) 5 units
 - (b) 7 units
 - (c) 9 units
 - (d) none of the above
- 31. If $y = x^3 9x^2 + x 2$, then the value of x for which $\frac{d^2y}{dx^2} = 12$ is
 - (a) 3
 - (b) 6
 - (c) 5
 - (d) 1

- 32. If $U = x^3 2y^3 + 3x^2y$, then $x \frac{\partial U}{\partial x} + y \frac{\partial U}{\partial y}$ is
 - (a) 3
 - (b) (
 - (c) 3u
 - (d) 2u

33.
$$\int_{0}^{\log 2} \frac{e^{x}}{e^{x} + 1} dx =$$
(a) log4

- (b) 3 log 2
- (c) $\log (\frac{2}{3})$
- (d) $\log \left(\frac{3}{2}\right)$
- 34. The algebraic sum of deviations of 10 observations from the constant C is -14 and the A.M. of the observations is 7.6. The value of C is
 - (a) 8
 - (b) 6
 - (c) 9
 - (d) 5
- 35. If the H.M. of 4, x and 6 be 6, the value of x is
 - (a) 8
 - (b) 12
 - (c) 10
 - (d) 6
- 36. A.M. and G.M. of two numbers are 10 and 6 respectively. The numbers are
 - (a) 2 and 18
 - (b) 3 and 17
 - (c) 4 and 9
 - (d) none of the above
- 37. Heights (in cm) of 8 boys are: 70, 74, 70, 73, 72, 70, 74 and 70. The median is
 - (a) 70 cm
 - (b) 71 cm
 - (c) 72 cm
 - (d) none of the above
- 38. If $\sum_{i=1}^{4} x_i = 32$ and $\sum_{i=1}^{4} (x_i 3)^2 = 120$, then $\sum_{i=1}^{4} x_i^2$ is
 - (a) 276
 - (b) 303
 - (c) 108
 - (d) none of the above

- 39. Relation between two variables x and v is 3x v + 11 = 0. If the median of y be 14, the median of x is (a) 1 (b) 2 (c) 3 (d) none of the above 40. The sum of 11 observations is 231 and mode is 18. The median of the observations is
- (a) 20

 - (b) 16
 - (c) 12
 - (d) none of the above
- 41. A.M. of n observations x_1, x_2, \dots, x_n is 15 and $\sum_{i=1}^{n} (x_i 9) = 72$.

The value of n is

- (a) 12
- (b) 16
- (c) 21
- (d) none of the above
- 42. The means of two groups of *n* and 12 observations are 33 and 40 respectively and the combined group mean is 36. The value of n is
 - (a)
 - 18 (b)
 - 14 (C)
 - (d) 16
- 43. If 3u = 2x and the H.M. of x is 0.09, the H.M. of u is
 - (a) 0.03
 - (b) 0.04
 - (C) 0.06
 - (d) none of the above
- 44. Daily wages (in ₹) of 5 workers are 90, 100, 70, 140 and 150. When the highest wage earner is absent on a particular day, the percentage change in range is
 - (a) 10.5
 - 12.5 (b)
 - 14.3 (c)
 - none of the above (d)

- 45. If $\sum_{i=1}^{10} X_i = 120$ and $\sum_{i=1}^{10} X_i^2 = 1690$, the standard deviation is

 - (b) 5
 - (c) 6
- The relation between two variables x and y is 2x + 5y =15. If the mean deviation of x about its mean 5 be 4, the mean deviation of y about its mean, is
 - (a) 1.4
 - (b) 1.8
 - (c) 1.2
 - (d) 1.6
- 47. The cost of manufacturing an article is ₹ 180 and it is represented in a pie chart. If electricity cost is given by 100°, the sum spent on other heads is
 - (a) ₹130
 - (b) ₹140
 - (c) ₹150
 - (d) none of the above
- 48. 50 workers in a plant earn ₹ 25,000 monthly with variance of wage distribution as 100. The coefficient of variation for plant is
 - (a) 1%
 - (b) 2%
 - (c) 2.5%
 - (d) none of the above
- For a moderately skewed distribution, mean = 20, coefficient of skewness = 0.25 and coefficient of variation is 20%. The mode is
 - (a) 19
 - (b) 17
 - (c) 12
 - (d) 15
- For a shoe-maker, which one of the following measures is most suitable to decide on the shoe sizes to manufacture?
 - (a) Median
 - (b) Arithmetic mean
 - Mode (c)
 - Geometric mean

Answer:

- 1. (c) 105:140:168:192
- 2. (a) 190 litre
- 3. (c) 10 days
- 4. (d) ₹1543
- 5. (b) 8 months
- 6. (c) 4
- 7. (c) 4
- 8. (a) 0
- 9. (c) 4
- **10.** (a) Φ
- 11. (c) (n! + 1):(n!-1)
- 12. (c) 9.5
- 13. (c) ~(~ p _v ~q)
- 14. (b) 54 sq cm
- 15. (d) none of the above
- 16. (a) 10 cm
- 17. (d) 15 cm
- 18. (b) 12 cm
- 19. (d) none of the above
- 20. (b) 12 cu cm
- 21. (c) 40√39sq cm
- 22. (d) None of the above
- **23.** (d) $\frac{\sqrt{5}}{2}$ units
- 24. (b) 4
- 25. (c) 6 units
- 26. (a) (-4, 1)
- 27. (d) None of the above
- 28. (b) 1
- **29.** (d) $\frac{\sqrt{x}}{2}$
- 30. (b) 7 units
- 31. (c) 5
- 32. (c) 3u
- 33. (d) $\log (\frac{3}{2})$
- 34. (c) 9
- 35. (b) 12
- 36. (a) 2 and 18
- 37. (b) 71 cm
- 38. (a) 276
- 39. (a) 1
- 40. (a) 20
- 41. (a) 12
- 42. (d) 16
- 43. (c) 0.06
- 44. (b) 12.5
- 45. (b) 5
- 46. (d) 1.6
- 47. (a) ₹130
- 48. (b) 2%
- 49. (a) 19
- 50. (c) Mode